




PATENT APPLICATION SERIAL NO. \_\_\_\_\_

71634 U.S. PTO  
60/043683  
04/14/97

U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE  
FEE RECORD SHEET





71634 U.S. PTO  
60/04



PTO/SB/16 (12-96)

Not for use through \*\* > 01/31/98 <. OMB 0451-001

Patent and Trademark Office U.S. DEPARTMENT OF COMMERCE

**PROVISIONAL APPLICATION >FOR PATENT< COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION >FOR PATENT< under 37 CFR 1.53 (b)(2).

Doclet Number		Type a plus sign (+) inside this box → +	
INVENTOR(S)/APPLICANT(S)			
LAST NAME	FIRST NAME	MIDDLE >NAME< INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Stapleton	Marilyn	J.	Durham, North Carolina NC
Sundseth	Rebecca		Durham, North Carolina NC
TITLE OF THE INVENTION (250 characters max)			
Immobilizing and Processing Specimens on Matrix Materials For The Identification of Nucleic Acid Sequences			
CORRESPONDENCE ADDRESS >(including country if not United States)<			
205 Winterberry Ridge Drive Durham, North Carolina 27713			
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification	Number of Pages	31	<input checked="" type="checkbox"/> Small Entity Statement
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	10	<input type="checkbox"/> Other (specify) _____
METHOD OF PAYMENT (check one)			
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fee			PROVISIONAL FILING FEE AMOUNT (\$) \$75.00
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fee and credit Deposit Account Number: _____			

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME

*Marilyn J. Stapleton*  
*Rebecca Sundseth*  
MARILYN J. STAPLETON  
REBECCA SUNDSETH

Date 4/14/97

REGISTRATION NO. \_\_\_\_\_  
(if appropriate)

☐ Additional inventors are being named on separately numbered sheets attached hereto

**>USE ONLY FOR FILING A< PROVISIONAL APPLICATION \*\*>FOR PATENT<**

Burden Hour Statement: This form is estimated to take 2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the \*\*> Chief Information Officer <, Patent and Trademark Office, Washington, DC 20231. \*\*DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.



PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IMMOBILIZING AND PROCESSING SPECIMENS ON MATRIX MATERIALS FOR  
THE IDENTIFICATION OF NUCLEIC ACID SEQUENCES

Inventors: Marilyn J. Stapleton and Rebecca Sundseth,

Express Mail mailing label no. EH949879765 US  
Date of Deposit 4-14-97  
I hereby certify that this paper or fee is being deposited with the United  
States Postal Service "Express Mail Post Office to Addressee" service  
under 37 CFR 1.10 on the date indicated above and is addressed to the  
Commissioner of Patents and Trademarks, Assistant Commissioner for  
Patents, Washington, D. C. 20231

Marilyn J. Stapleton  
Typed or printed name of person mailing paper or fee

Marilyn J. Stapleton  
Signature of person mailing label or fee

Please accept provisional application for patents as titled above and enclosed herein.  
A check for \$75.00 is enclosed.

#10

75<sup>00</sup>-214

A1 PROVISIONAL

1

IMMOBILIZING AND PROCESSING SPECIMENS ON MATRIX MATERIALS FOR  
THE IDENTIFICATION OF NUCLEIC ACID SEQUENCES

BACKGROUND OF THE INVENTION

This application is a continuation-in-part application of U. S. Patent Application Serial No. 935,637, now U. S. Patent No. 5,382,511 and U. S. Patent Application Serial No. 836,348, now U. S. Patent No. 5,451,500.

FIELD OF THE INVENTION

The present invention relates to a process and device for analyzing a biological specimen for the presence of a nucleic acid sequence for diagnostic purposes. More specifically, it relates to immobilizing small amounts of cellular or tissue specimens for the analysis of nucleic acids without extracting them from the specimen. Immobilizing cells and tissue simplifies sample preparation for a variety of different nucleic acid detection methods. In some instances the number of copies of the nucleic acid sequence of interest present in the specimen is sufficient for detection, particularly with newer, more sensitive detection methods. In other instances, the nucleic acid sequence of interest is extensive enough that detection is possible by increased labeling of the target or probe molecules. In other instances when the nucleic acid sequence of interest is rare in the specimen or when distinguishing fine sequence detail, a means to amplify the nucleic acid sequence

above background may be desirable for detection. The process and device of the invention utilizes immobilization of the biological specimen to a matrix for the purpose of detecting specific nucleic acid sequences in the specimen. Detection occurs when either a reagent that recognizes the target nucleic acid binds to targets within the biological specimen and the reagent's presence is detected in the matrix, or a first reagent modifies a specific target sequence of the specimen in a way that the modified product sequence is detected by the location where it binds to a second reagent. Means to modify a target sequence include enzymatic means to synthesize a nucleic acid sequence from one or more specific primers or ligate two sequences together based on their binding adjacently on a specific template. Means to detect the modified target sequences include binding to a second reagent that recognizes the modified product and is in a specific location, such as a probe array. Immobilized Sample Amplification, or ISA, increases the labeled molecules or number of copies of the molecular target adequately enough for detection by appropriate means. Appropriate means include, but are not limited to, detection of signals that are fluorescent, luminescent, colorimetric, electrical, chemical or image analysis and any combinations thereof.

#### DESCRIPTION OF RELATED DISCLOSURES

In U. S. Patent Number 5,382,511, a process to amplify nucleic acid targets was invented wherein a biological specimen was immobilized on the surface of a microscope slide or embedded in a gel matrix, for preparing and directly amplifying the nucleic acids in

the sample. Prior to said method invention, nucleic acid amplification had been limited to solution reactions wherein the nucleic acids were first extracted from cells or tissue. A number of examples for using *in situ* amplification are given in U. S. Patent Number 5,188,963. A photomicrograph of cells which had amplified and labeled DNA was included in U. S. Patent Number 5,451,500 to show that the amplified fragments are retained in individual cells and such cells can be enumerated under microscopic observation. A number of subsequent publications support this method and even the localization of nucleic acid amplification products on cells treated in a manner in which only the chromosomes remain immobilized on the glass slides.

In some nucleic acid diagnostic tests, microscopic observation of the biological specimen to look for morphological changes is productive and informative to a diagnosis. The degree of cell fixation needed to maintain cell morphology may be counterproductive to efficient detection or amplification because the most common cell fixatives cross-link the proteins and nucleic acids. Protease digestion is frequently used to reverse crosslinking because crosslinks interfere with hybridization to target sequences or terminates DNA primer extensions during DNA synthesis, thereby reducing the efficiency of amplification or hybridization detection.

In many DNA-based diagnostic tests, maintaining cellular morphology may be irrelevant to a diagnosis as long as the origin of the specimen is known. For example, the specimen may be aspirated from a series of a patient's lymph nodes, so that a pattern of metastasis from a primary tumor may be identified by the presence or absence of specific nucleic acid tumor markers in each of the lymph nodes.

The standard research protocol for sample preparation is to lyse cells to release the nucleic acids. Nucleic acids are then separated from the cell's peptides and lipids by phenol:chloroform extraction and then ethanol-precipitated to concentrate them. Varying amounts of the sample's nucleic acids are lost during the many extraction steps. Special DNA purification materials are available, but require filtration, centrifugation, or electrophoresis and add cost to preparation.

The conventional approach for clinical specimens is to perturb the cells or viral particles from a few cells or a small amount of tissue in order to release nucleic acids into solution and use a portion of the crude extraction. When this approach fails, the nucleic acid component of the crude extract requires further purification. A variety of methods exist to enzymatically either label or amplify specific nucleic acid sequences in order that they may be detected. Amplifying DNA or RNA from clinical specimens has been difficult to optimize because the clinical specimens present so many different parameters and potential inhibitors. Quick lysis techniques give rise to unpredictable or anemic PCR reactions. Anemic reactions are a continual challenge in making a robust diagnostic test because detection depends upon reproducibly amplifying the specific targets from a complex genetic background.

The *in situ* amplification process described in U.S. Patent Numbers 5,382,511 and 5,451,500 uses enzymes such as polymerase or ligase, separately or in combination, to repeatedly generate more copies of a target nucleic acid sequence by primer extensions to incorporate new nucleotides or by ligations of adjacent complementary oligonucleotides, wherein each template generates more copies and the copies may themselves become template. By melting complementary strands of nucleic acids, the original strand and each

new strand synthesized are potential templates for repeated primer annealing or ligation reactions to make and expand the number of specific, amplified products. A thermostable polymerase with reverse transcriptase activity and a thermostable ligase are now both commercially available and increase the choice of enzymes for both RNA and DNA detections. The amplification can either be primer extensions in one direction for linear amplification, or in opposing directions, for geometric amplification. The label can either be incorporated as labeled nucleotides or labeled primers for one-step detection or labeled probes may be added whereby the probes hybridize to the amplified products for detection. Other schemes for modifying, labeling or amplifying, include, but are not limited to, PCR (Roche Molecular Systems, Branchburg, New Jersey), NASBA, (Organon Teknika, Durham, North Carolina) or SDA, Becton Dickinson, (Franklin Lakes, New Jersey), assays in which peptide nucleic acids recognize and bind target sequences and assays in which reagents recognize mismatches between the specimen's nucleic acid sequence and a known standard. These are variations of modifying or making multiple copies of specific nucleic acid templates for the purpose of detection, and when performed directly on an immobilized biological specimen are within the scope of either this invention or the heretofore mentioned U.S. Patent Numbers 5,382,511 and 5,451,500.

Microporous matrices are generally used in the field of molecular biology as a solid support media for proteins or nucleic acids transferred from electrophoretic gels. The Southern transfer technique is used to blot nucleic acids from gels onto a matrix in order to detect specific sequences by hybridization. In other instances, nucleic acids are filtered or spotted directly onto the matrix. The mechanism of the binding of proteins or DNA to matrices depends mainly on a combination of hydrophobic interactions, hydrogen bonding,



and salt bridges. Nitrocellulose matrices were the first type of media used for Southern transfer and required baking in a vacuum oven to make the bond formed between the medium and the nucleic acid essentially permanent, permitting stringent assay conditions and reuse of the blot after removal of a bound probe. One criteria for solid supports is that all reactive sites on the surface surrounding the bound molecule be eliminated or rendered inert to non-specific interactions. Properties of different types of matrices made of diazotized media, nylon and ion exchange matrices are matched to different applications and the size of the molecules to be retained. DEAE anion exchange matrices are used for reversible transfer-elution to isolate RNA or DNA wherein different buffers favor the binding or release of the biomolecules to the medium.

More hydrophilic matrices have been formulated for better wetting and less flow resistance. Manufacturers have developed blood separation media to trap white blood cells preferentially. Another fibrous medium separates the cellular and noncellular fractions of whole blood, as it wicks through the porous spaces in the matrix. The chromatographic effect that occurs during wetting with a given volume of whole blood displays a constant ratio of 25% for the cell-free plasma fraction in the Hemadyne matrix, Pall Corporation, Port Washington, New York. The pore structure of the matrix and its low binding affinity allows even high molecular weight proteins to migrate unimpeded and separate in an aqueous phase apart from the blood cells. These matrices present a quick way to either filter white cells from blood or separate biomolecules in the plasma away from the red blood cells. The manufacturer recommends extracting the nucleic acids from the matrix by using a proteinase enzyme to free the DNA from the immobilized white blood cells. The cell lysate, or a more purified nucleic acid portion, is then added to a

polymerase chain reaction (PCR). Separating white cells or biomolecules away from the red cells is useful in sample preparation because the large amount of hemoglobin in blood is known to poison enzymatic reactions such as PCR.

#### SUMMARY OF THE INVENTION

The method and device of this invention minimizes the quantity of a specimen required for a diagnostic result because the nucleic acids present in an immobilized specimen are labeled or amplified directly without the losses that are associated with extracting them from the specimen. The present invention immobilizes a small amount of biological or clinical specimen such as blood, needle biopsy aspirate, cell smear, cell print or ultra-thin tissue specimen onto a supporting matrix material, and dehydrates the specimen quickly to prevent degradation of nucleic acids. The matrix material containing the immobilized specimen is then rehydrated with either a labeling or amplifying reaction mixture for the detection of nucleic acids sequences of diagnostic interest.

The present invention is useful in reducing the quantity of reagents needed for processing the specimen. Fibrous materials increase the surface area relative to volume over solid supports. Reagents fill the void volume of a fibrous matrix and readily access the target nucleic acids expected to be present in an immobilized biological sample. Sample preparation is therefore miniaturized and simplified by processing the sample immobilized on a fibrous matrix. The matrix material to which the sample is dried may be rinsed or pretreated to remove any interfering components of the specimen and dehydrated a second time before detection.

The present invention improves the containment and integrity of patient specimens by collecting and transporting the specimen via the matrix material. The specimen is dried onto the solid-support matrix and amplified without using any of the standard methods to remove nucleic acids from cells or viral particles. The immobilized specimen, and the nucleic acids present, are treated less harshly than standard treatments with organic solvents. There is no need to recover and transfer nucleic acids from one reaction vessel or mixture to another. More specifically, the present invention enables automated processing by integrating the sample preparation with an enzymatic labeling or amplification step.

In accordance with the present invention, the preferred embodiment of the device comprises a fibrous matrix that is of a thin, flat shape rather than spherical or cuboidal, wherein a biological specimen is brought into contact with the total surface area of the fibrous matrix and then dehydrated by evaporation. A preferred embodiment of the size and density of matrix fibers is dependent upon the type of specimen.

In yet another embodiment, the invention herein relates to a device comprising a fibrous matrix attached to a handle for the purpose of touching the matrix to biological tissue. The device collects a cellular specimen more easily than trying to hold the tissue specimen and to touch it to a matrix or glass slide because the smaller matrix surface can be directed to a specific area of the specimen.

In a broad aspect, the component device of the invention comprises means for immobilizing a biological specimen by first drying it to a matrix support and then filling the void volume of the matrix with liquid reagents to amplify, label or hybridize specific

nucleic acids for detection. The invention is readily adaptable for use in an automated analyzer for DNA diagnostic tests.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1A and 1B illustrate genetic products amplified from human blood immobilized and dried on different matrix materials and which are visualized after electrophoresis in agarose gels and showing bands the size of the expected targets.

FIGURE 2A and 2B illustrate genetic products of viral amplification from immobilized cell culture supernatant, whole blood, and plasma which was fractionated from whole blood by lateral flow through Hemadyne matrix material before immobilization, and visualized after electrophoresis in agarose gels.

FIGURE 3A and 3B illustrate agarose gel electrophoresis of products, amplified from equine dermal cells that were lifted from glass and plastic culture dishes by contact with a matrix material, immobilized and processed on the matrix material.

FIGURE 4 illustrates agarose gel electrophoresis products amplified from human cheek cells collected, immobilized and processed on a matrix material.

FIGURE 5 illustrates agarose gel electrophoresis products amplified from serial dilutions of whole blood immobilized on a matrix from as few as 100 white cells.

FIGURE 6 illustrates agarose gel electrophoresis products amplified from mRNA, as well as DNA, present in human cells immobilized and processed on a matrix material.

FIGURE 7 is a drawing of an embodiment of a sample collecting device for the analysis of nucleic acids.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention broadly comprises a method to immobilize biological and cellular specimens on a matrix for detection of target nucleic acids in the specimens. In accordance with the present invention, the preferred embodiment of the device comprises a matrix that is of a thin, flat shape rather than spherical or cuboidal, wherein a biological specimen is brought into contact with the surface area of the matrix and then dehydrated by evaporation. Dehydration is more rapid if the matrix is thin. Dehydrating is significant because the volume of the original sample may be larger, relative to the reaction volume, than if the sample was introduced as a liquid into the same reaction volume. In other words, dehydration is a way to concentrate the cells or virions containing nucleic acids from an aqueous, liquid biological sample and process more nucleic acid analytes in a smaller reaction volume and in a smaller space. A membranous matrix comprising more thread-like fibers and smaller fiber diameters increases the surface area of binding sites to immobilize all of the different components of a biological specimen. Additionally, a surface area with more binding sites than needed to adhere just the cells also serves to bind and effectively sequester those molecules which may inhibit nucleic acid modifying and amplifying enzymes.

The capillary air spaces in a fibrous matrix causes liquids and soft tissue cells to flow and spread throughout the matrix. This wicking or absorbing property of fibrous matrix materials causes more cells to adhere to a fibrous matrix having an outer dimension equivalent to an area of a smooth, flat glass slide. Different biological tissues vary

considerably in cell density and in the viscosity of their intercellular matrix material. A fluid tissue like whole blood is expected to wick further into the void spaces of a matrix than a tissue such as adenoids that are less fluid. A matrix comprised of a low density of fibers will contain a larger specimen volume than a high density matrix with a lesser void volume. The physical dimensions and composition of the matrix are selected on the basis of the general size and viscosity of the specimen so that the density of its fibers is sufficient to absorb the liquid volume of the specimen and immobilize the cellular components.

Individual cell clusters, or cells comprising desired characteristics, may be identified visually or microscopically and collected by physical contact with a fibrous matrix. The matrix containing the selected cellular specimen may be dried and analyzed for specific DNA and RNA content without isolating nucleic acids from the cellular material. A 3 x 3 millimeter square area may immobilize upwards of 200,000 cells from a biological specimen; however the sample may only need to contain ten to twenty cells or gene targets for amplification in order to increase the number of target copies to a detectable level. The capacity of a fibrous matrix to immobilize cells, is a desirable feature to incorporate into a device for analyzing genetic targets.

The number of cells needed in a diagnostic sample is determined statistically based on whether the target nucleic acid sequences occur commonly or rarely in a cell population. A fibrous matrix having one-eighth inch sides (3.2 millimeters) and a depth of 0.020 to 0.040 inch (0.5 - 1 millimeter) that is filled with a fluid layer has a volume equivalent to 5 - 10 microliters less the space filled by the fibers. Reducing the total reaction volume to a range of 5-25 microliters will conserve reagents and lower costs.

In cases where nearly all the cells in a specimen are expected to contain the target sequence, only a few cells are needed in order to amplify genetic sequences present even at only the single copy level per cell. The specimen surface area needed for a few cells is in the range of less than 100 square microns and the volume of reagents is in the range of a few picoliters. In cases where the target is expected to be a rare event, the specimen surface area may be accordingly increased in order to contain around 25 microliters of reagents for thousands of cells. Liquid droplets of cells dry onto a planar surface with cells affixed generally in a monolayer, if the cells are not too concentrated in the solution. Embedded or frozen tissue is routinely cut into 4-10 microns sections that are nearly a monolayer or at most a few cells thick. With the device of the invention, cells are collected and immobilized similarly in a monolayer-like arrangement, but by means that do not involve embedding in a liquid media phase that solidifies for microtome sectioning. The immobilized, dehydrated specimen preserves labile mRNA and is directly accessible for the diffusion of amplifying and hybridizing reagents.

The dimensions of small fibrous matrix pieces, and a device for handling and enclosing them in a reaction chamber, will depend upon the density of cells and other biomolecules in the specimen. It is expected that specimens containing more cells and biomolecules will occupy more surface area of the matrix, depending on their ability to flow. The reaction volume needs to provide sufficient reactants so as to not limit biochemical activity. The surface area of the fibrous matrix used should not be in large excess of what is needed to immobilize the specimen since the remaining reactive sites on the matrix may bind the amplification enzymes, primers and nucleotides, labeled or not, limiting their availability for the amplification or detection reaction.

Chambers, as described in applicant's U. S. Pat. No. 5,346,672, and having a specimen area capacity of a 15 millimeter-diameter circle on a microscope slide are routinely used in our laboratory to prevent evaporation during thermocycling slide specimens. As a matter of standard procedure, we removed the reaction liquid after amplification and analyzed it by agarose gel electrophoresis because products from the slide specimen were found to be present in the solution. Results were used to confirm the size of products amplified from the specimen immobilized on the slide. Frequently, the slide specimen area will contain 50,000 cells or less and cover only a 3 millimeter diameter area on the slide surface. The chamber contains a liquid volume of approximately 50 to 75 microliter. Under conditions when the cells are covering only a portion of the positively-charged surface of an adherent slide enclosed in the reaction chamber, it is necessary to include in the amplification mix 0.1-0.01% bovine serum albumin (BSA), or a similar blocking agent, in order to bind the unoccupied plus charges on the slide and block the polymerase enzyme and nucleic acid molecules from binding to the slide. The amount of BSA needed for an optimum amplification reaction may be empirically determined. Using 0.01% BSA in immobilized sample amplifications has become routine as a precaution to cover reactive sites on matrix materials that may interfere with enzyme activity.

The applicants performed hundreds of genetic detection reactions on biological specimens, namely blood and cheek cells. Fresh whole blood from fingerpricks and cheek cells collected by either whole mouth fluids, transmucosal exudate and buccal smears, were immobilized on a dozen of different matrix materials from several manufacturers (namely, Gelman Sciences, Ann Arbor, MI; Magna Separations Inc. (MSI), Westboro, MA; Pall BioSupport, Port Washington, NY; Porex Technologies, Fairburn, GA;



Schleicher & Schuell, Keene, NH; Tetko, Briarcliff Manor, NY; 3M, St. Paul, MN). It was convenient in the laboratory to detect by amplifying and subjecting the product and controls to agarose gel electrophoresis and UV illumination of ethidium bromide fluorescence. However, it is understood there are many other ways known in molecular biology to detect nucleic acids, many of which detect fewer molecules than can be visualized by the detection method used herein.

#### EXAMPLE I

Early experiments in our laboratory confirmed that ISA products from whole blood spotted on matrices were the same as the products obtained from amplifying purified human DNA spotted on the same type of matrix. Bands in Figure 1A demonstrate the same specific 524 base-pair fragment, amplified by using primer extensions from two opposing primers complementary to a region of the human p53 gene. 100  $\mu$ l reactions were comprised of 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 0.01% BSA, 0.2 mM of each of the 4 standard dNTPs, 1.5 mM MgCl<sub>2</sub>, 0.1  $\mu$ M primers and 1.25 Units of Taq DNA polymerase. Thermocycling conditions were an initial denaturation at 94 °C for 4 min., followed by 30 cycles of 94 °C, 45"; 52 °C, 45"; 72 °C, 1' and then a final extension at 72 °C for 5'. Following thermocycling, twenty microliters of each reaction were electrophoresed on a 1.5% agarose gel and products visualized with UV-illumination of the ethidium bromide stained gel. Lanes 2 and 3, represent ISA product from five-microliter, dried whole blood spots on Hemadyne, rinsed briefly with water and 1 mM HCl, respectively, and lane 4, product from a dried spot of 100 nanograms of human genomic DNA. (Lane 1 is Boehringer Mannheim DNA molecular Marker VI).

Reports by others in the literature that the presence of hemoglobin inhibits enzymatic amplification apparently is minimized when the blood is immobilized on a matrix. Further experiments compared immobilizing blood on a variety of different matrix materials. Fresh whole human blood was collected directly from a finger prick and 2  $\mu$ l was spotted onto each of several small squares of different matrix materials. The sizes of the matrices were approximately 4-9 mm<sup>2</sup>. Within seconds the blood flowed into the intramatrix spaces and in a few minutes the blood components dried to the surface of the matrix fiber material. For the inactivation of blood borne pathogens, the matrix materials containing blood are heated at 82 °C for 15 min. Each matrix was washed twice for 5 min each with 100  $\mu$ l of water and redried at 82 °C for 15 min. The matrices containing the cells were then amplified in thin-walled tubes under the following conditions in 50  $\mu$ l reactions: 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 0.01% BSA, 0.2 mM of each of the 4 standard dNTPs, 2.25 mM MgCl<sub>2</sub>, 0.5  $\mu$ M primers and 1.25 Units of Taq DNA polymerase. Thermocycling conditions were an initial denaturation at 94 °C for 3'50", followed by 30 cycles of 94 °C, 45"; 60 °C, 45"; 72 °C, 1' and then a final extension at 72 °C for 5'. Following thermocycling, twenty microliters of each reaction were electrophoresed on a 3% agarose gel and products visualized with UV-illumination of the ethidium bromide stained gel. Primers, used for identity testing and recognizing a single copy locus (locus 34), generated the appropriate 115 base-pair amplification product from blood samples immobilized on the following matrices as shown in Figure 1B: lane 2, Nylon 3-3710, (Tetko); lane 3, Nitex Nylon (Tetko); lane 4, Magna nylon (MSI); lane 5, Hemadyne (Pall); lane 6, Leukosorb (Pall); lane 8, polyester twill (Tetko); lane 9, PeCap polyester 17/9 (Tetko); lane 10, PeCap polyester 43/29

(Tetko); lane 11, PeCap polyester 150/41 (Tetko); lane 12, PeCap polyester 210/35 (Tetko); lane 13, Tetex polyester (Tetko). Lane 1 represents a control reaction containing 100 ng purified human genomic DNA and Lane 14 is Boehringer Mannheim DNA molecular Marker XIII. There was no band apparent in lane 7, representing NFWA matrix (Gelman). The invention is a method for the analysis of nucleic acids of a biological specimen, comprising (a) immobilizing a given specimen on the surfaces of a fibrous matrix material, (b) amplifying a targeted sequence region of said specimen's nucleic acid, and (c) detecting the presence or absence of said targeted sequence of said specimen's nucleic acid. Accordingly, the invention is a method for the analysis of nucleic acids wherein the biological specimen is blood.

## EXAMPLE 2

Hemadyne lateral flow membrane was used to separate whole blood into a cellular fraction and a cell-free plasma fraction in order to investigate the capability of the RNA retrovirus, Equine Infectious Anemia Virus (EIAV) to flow through the membrane with a plasma fraction and to be identified by ISA. One microliter of the supernatant of EIAV-infected equine dermal cell culture, was spotted onto Hemadyne alone or mixed with 5 microliters of human whole blood first and then spotted on Hemadyne, as well as whole blood alone. Sections of the Hemadyne membranes containing either the dried culture supernatant or plasma, which had been separated from the cellular fraction, were excised and used for ISA. The matrix pieces were added to separate reverse transcription reactions using SuperScriptII Reverse Transcriptase (Life Technologies, Gaithersburg,

MD) according to manufacturer's instructions and incorporating an EIAV-specific primer. One-fourth of each reverse transcription reaction was amplified by Taq polymerase with an EIAV primer pair. In Figure 2A, bands representing product amplified from EIAV virus immobilized on the Hemadyne were visualized in all lanes except lanes 5 and 6 in which no EIAV had been added to the blood; lane 2, unwashed, dried culture supernatant spot; lanes 3 and 4, washed, dried culture supernatant spot; lanes 7 and 8, EIAV in the plasma fraction. The invention is a method of detecting viral nucleic acids present in a whole blood specimen, comprising (a) applying said whole blood specimen to a lateral flow matrix; (b) separating said whole blood specimen into a cellular fraction on a first portion of said lateral flow matrix and a plasma fraction on a second portion of said lateral flow matrix; (c) drying and immobilizing said plasma fraction on a second portion of said lateral flow matrix; (d) saturating said plasma fraction on a second portion of said lateral flow matrix with amplification reagents to enzymatically amplify viral nucleic acid targets present in said plasma fraction; (e) determining whether said plasma fraction of said whole blood specimen contains said viral nucleic acid targets.

In other experiments, results shown Figure 2B, 10 microliters of horse blood, with EIAV-containing supernatant added (lanes 6-9), and without EIAV (lanes 2-5), was applied to one end of a Hemadyne strip and flowed laterally across the strip. After lateral flow had ceased, the Hemadyne strip was dried and then cut into four sections and each section was amplified using rTth DNA polymerase (Perkin Elmer, Foster City, CA), EIAV primers and appropriate buffers and thermocycling conditions. The blood that was pipetted onto a portion of the matrix strip is represented in lanes 2 and 6; and flowed laterally through portions of the matrix represented in lanes 3-4 and 7-8, respectively, and

portions of the matrix containing plasma where the flow ended are represented in lanes 5 and 9. The bands in Lanes 6-9 are evidence that viral particles are deposited over the entire area of the matrix. The blood was taken from a horse previously infected with EIAV and expected to have very low amounts of virus remaining. The bands in lanes 6-9 and a faint band in Lane 1 (visible on the original photograph) are evidence that ISA is capable of detecting virus in blood when either sufficient virus present, the volume of blood sample is adequate or a method more sensitive than an agarose gel is used to detect amplified product. When the virus was present at very low copy number, as in the infected horse blood, a band was seen after amplification only in the portion of the matrix where blood was deposited directly on the membrane; when excess virus was added to the blood, the virus was detectable in all portions of the lateral flow matrix. In experiments performed at the veterinary school from where the horse's blood was obtained, extensive measures are required to amplify detectable EIAV from this same horse by standard means. First white cells are isolated by a Ficoll gradient, nucleic acids are purified from the white cells, and two sequential PCR are performed with a portion of the first reaction mix being used in the second reaction. Viral RNA or DNA is therefore available more readily for enzymatic amplification from whole blood specimens and component fractions of whole blood by the ISA method than standard PCR methods.

The invention is a process of detecting viral nucleic acids present in whole blood, comprising (a) applying a specimen of whole blood to a fibrous matrix; (b) drying said specimen of whole blood on the surfaces of said fibrous matrix; (c) applying reagents to said fibrous matrix to enzymatically amplify viral nucleic acid targets that may be present in said specimen of whole blood; (d) determining whether said whole blood specimen

contains said viral nucleic acid targets. Accordingly, by comparing results to known quantities of virus and blood volumes, the relative quantities of said viral nucleic acid targets present in said specimen may be measured. A standardized amount of a mimic virus, one constructed to have a sequence variation to make it detectable, as well as complementary sequence structure so that it would be amplified in the same manner, may be added to and immobilized together with the specimen in order establish a positive control for the reaction and quantify relative amounts of virus particle present in the specimen. The invention is a method of detecting viral nucleic acids further comprising means to determine quantities of said viral nucleic acid targets present in said specimen.

### EXAMPLE 3

Evidence for the ability of a fibrous matrix material to actively pick up and immobilize cells for detection of a gene sequence present at one or two copies per cell is shown in Figure 3. Equine dermal cells that had been grown to a confluent monolayer on either plastic tissue culture plates, or on sterile glass chamber slides, were washed twice with phosphate buffered saline and drained briefly to remove excess PBS while also keeping the cells adhered to the plastic or glass. Small squares of matrix (approximately 4-9 mm<sup>2</sup>) were gently touched to the surface of a portion of the moist cell monolayer to collect cells. The matrices were then transferred to reaction vessels and dried and washed as described above. Reaction mixtures were as described for Figure 1B, except for a change to 1 mM MgCl<sub>2</sub> and primers recognizing single-copy, equine gene sequences for tumor necrosis factor and a 55 °C

annealing temperature were used. (Lane 1 contains Boehringer Mannheim molecular weight marker XIII.) The expected 245 base-pair amplification product was obtained in Figure 3, lanes 2 and 3 from horse genomic DNA and also from cells grown on either plastic (lanes 4-9) or glass (lanes 10-15) and printed onto one of three matrices for immobilized sample amplification. (Nylon 3-3710, lanes 4,5,10,11; PeCap polyester 17/9: lanes 6,7,12,13; PeCap polyester 43/29: lanes 8,9,14,15).

Small pieces of matrices are difficult to handle whether manufacturing or using them as devices. Squares cut from Nylon 3-3710 were adhered with a double-sided adhesive transfer tape to a polycarbonate film of the same length and width of the matrix square, but having a thickness of 0.007 inch (178 microns), to support the matrix and make it more easily handled. The results of Figure 3B, lanes 5 and 7 show that the matrix's ability to pick up cells and amplify nucleic acids from them, under conditions as described in Figure 3A, was not altered or compromised when a piece of the matrix was attached to an adhesive and backing for support. (Figure 3B, lanes 2 and 3 contained products amplified from horse genomic DNA and lanes 4 and 6 contain products on unsupported pieces of matrix, touched to cells on plastic and glass culture dishes, respectively, as in Figure 3A.) The sandwiched device comprising a matrix, adhesive and backing is in communication with enzymatic amplification and hybridization reagents; therefore all materials used in fabricating different embodiments of the invention would be tested for biocompatibility. The applicants have had experience with many different materials with which to fabricate devices and have identified some that do and some that do not affect ISA for the detection of nucleic acids. The invention is a device for analyzing the genetic sequence of small

amounts of the nucleic acids of a specimen, comprising (a) a fibrous matrix, (b) means to support said fibrous matrix, (c) means to contact said fibrous matrix with a biological specimen, (d) means for cellular material of said biological specimen to attach to said fibrous matrix, and (e) means to detect the presence or absence of specific genetic sequences in the nucleic acids of said cellular material transferred to said fibrous matrix from said biological specimen.

#### EXAMPLE 4

In another experiment, a piece of matrix was attached to the end of a handle with an adhesive that had been shown previously to be compatible with ISA. The unit was autoclaved and used to collect cheek cells directly from the mouth of a volunteer. After collection, each matrix with adhesive backing was detached from the handle and transferred to reaction vessels, dried and processed as described for Figure 1B. The matrices were rehydrated with a reaction mix containing the components and products were visualized as described previously for Figure 1B. In Figure 4, primers recognizing a single copy, identity-testing locus (locus 60) generated a 169 base-pair amplification product both in control reactions containing human genomic DNA target (lanes 2 and 3) and in ISA reactions containing cheek tissue cells immobilized on Nylon 3710, (lanes 5 and 6). No product was evident in reactions containing no target (lane 4). (Lane 1 contains Boehringer Mannheim DNA molecular weight marker XIII.) The invention is a device for analyzing the genetic sequence of small amounts of the nucleic acids of a human subject, comprising (a) a fibrous matrix, (b)



means to support said fibrous matrix, (c) means to contact said fibrous matrix with a human subject to obtain a biological specimen, (d) means for cellular material of said biological specimen of said human subject to attach to said fibrous matrix, and (e) means to detect the presence or absence of specific genetic sequences in the nucleic acids of said cellular material transferred to said fibrous matrix from said biological specimen of said human subject.

#### EXAMPLE 5

Biological specimens were diluted to derive the sensitivity level, or copy number, at which a particular nucleic acid sequence could be detected. The number of cells containing nucleic acids in human blood is relatively constant so blood was diluted and each dilution was analyzed for the detection of a genetic sequence represented with two alleles per cell. Fresh human blood was collected from two individuals by fingerprick and either spotted directly onto a matrix or diluted in phosphate buffered saline and then spotted directly onto a matrix. Each dried matrix was added to a reaction vessel with a polymerase chain reaction mix containing the components described for Figure 1B, with a substitution of 1.5 mM  $MgCl_2$ . Reactions were thermocycled using the same primers described for Figure 1B. In Figure 5, lanes 2 and 3 demonstrate the 115 base-pair product generated by specific amplification of human genomic DNA. The product amplified by ISA from 2  $\mu$ l of immobilized human blood can be seen in lanes 4 and 9. Two microliters of serial dilutions (1:5, lanes 5

and 10; 1:10, lanes 6 and 11; 1:50, lanes 7 and 12; 1:100, lanes 8 and 13) are also sufficient to generate a visible product on the original photograph of the gel. (Lane 1 contains Boehringer Mannheim DNA molecular weight marker XIII.) The results of Figure 5 demonstrate that a single-copy human gene can be amplified from 0.02  $\mu$ l of immobilized human blood, or approximately 100 cells. Bands were visible in other experiments from dilutions estimated to have as few as 20 immobilized cells.

Because immobilized sample amplification does not depend upon isolation techniques in which the recovery of nucleic acids may vary from sample to sample, rapid analysis of either single cells or a few cells immobilized on a matrix is more achievable. In certain research applications it may be desirable to analyze the nucleic acids of a single cell, but the interpretation of results for diagnostic purposes will be more reliable when based on more than a single cell. An advantage of ISA is the capability to analyze the genetic material of a small amount of biological specimen for a variety of different reasons: surgical biopsy material may be sparse as in needle aspirates of suspected breast tumors and lymph nodes; a small region may be selected to determine the extent or border of the malignant characteristics of a tumor by analyzing somatic genetic mutations; and regions of tissue may be analyzed after genetic therapy to determine the spatial or temporal distribution of the genetic intervention. Anatomical orifices and surface lesions are non-surgical sources of biological specimens available for ISA. For example, skin lesions caused by ticks are a more direct source of material to test for nucleic acid evidence of parasites than peripheral blood. Fresh cervical tissue from suspicious zones may be transferred by

touching to a matrix for the analysis of abnormal cytology, thereby eliminating scraping or incisions to do so.

In cases in which a few cells or viruses are contained in a larger volume of biological fluid than can conveniently be applied and dried to a small matrix, for example in the spinal fluid from lumbar punctures of patients with meningitis, the matrix itself may be used to capture the cells or viruses. Matrices have traditionally been used for separation and filtration by using the sieving mechanism of their pore size to separates biomolecules based on size. For ISA the capture does not need to be based on sieving but rather a natural affinity of cellular and viral surface biomolecules to attach to the matrix material. In our laboratory two different matrix materials were layered, each capable of binding cells, but the cells adhered to the matrix material it first contacted. The liquids continued to wick into the capillary spaces of the second matrix even though few cells were carried with it. It is obvious that for the purpose of keeping the matrix material small enough for a low volume reaction and minimizing the surface area of the matrix which enters the reaction, that the collecting device may be backed with an absorbent material which is not part of the matrix.

In many instances, fluids undergoing filtration are pressurized to force fluids through the matrix voids. Without pressurization, the capillary force of the aqueous liquids is the chief mechanism in filling the matrix voids, initially replacing the air in these spaces. Either using pressure, or only the capillary wicking action of the matrix, as a separation system is compatible with ISA. The fibrous and chemical nature of the matrices that comprise the sample collection device, and their ability to bind and immobilize the biological specimen containing genetic material to its surface area for the

analysis of nucleic acids, is enabling for the device of the invention. The device of the invention may optionally contain absorbent material in addition to the primary matrix.

#### EXAMPLE 6

The analysis of RNA expression from immobilized cells and RNA viruses is within the scope of the invention. In Figure 6 reverse transcription and amplification of messenger RNA was demonstrated in human cells immobilized on a matrix. A derivative of human lymphoblastoid CEM cells growing in suspension culture were concentrated by centrifugation and then resuspended in culture media containing serum. Aliquots containing approximately 10,000 cells were applied to small squares (4-9 mm<sup>2</sup>) of matrix, dried and washed as described above. The combined reverse transcriptase and amplification system sold by Boehringer Mannheim, as their Titan RT-PCR System, containing a 1x reaction buffer of 1.5 mM MgCl<sub>2</sub> and DMSO, 0.2 mM dNTP's, 0.01% bovine serum albumin, 5 mM dithiothreitol, 0.5 μM of each primer and 1 μl of Titan enzyme mix was added directly to the dried matrices. Reactions were subjected to an initial 30 minute incubation at 50 °C to allow reverse transcription of mRNA, followed by 30 cycles of 94 °C, 45"; 60 °C, 45"; 72 °C, 15" and a final extension incubation at 72 °C for 5 min. Oligonucleotide primers homologous to two exons of the tumor necrosis factor gene generate a 568 base-pair product of DNA amplification and a 266 base-pair product resulting from RT-PCR amplification of the spliced tumor necrosis factor messenger RNA, lanes 2 and 3 in Figure 6. The template for amplification in Lane 2 are purified nucleic acids isolated

from human CEM cells and in lane 3, the CEM cells immobilized on Nylon 3-3710 are amplified by ISA. In methods for amplifying RNA, other than the ISA method, strong denaturing solutions are added to cell lysates to preserve RNA. Extensive measures are taken to recover RNA rapidly before it is degraded. In other ISA experiments (data not shown), heating the immobilized cells and matrices in water for 20 minutes at 90 ° C. eliminated evidence of RNA amplification, but the capability to amplify DNA was unchanged. One advantage of ISA for mRNA detection is a simple means to detect mRNA directly from cells of interest without extraordinary measures.

The invention is a method for the analysis of a small amount of ribonucleic acid of a biological specimen, comprising: (a) immobilizing a biological specimen on surfaces of a fibrous matrix material; (b) amplifying a targeted region of said biological specimen's nucleic acid sequence; (c) means to detect the amplified targeted region of said biological specimen's nucleic acid sequence; in order to identify the presence or absence of at least one particular ribonucleic acid sequence in said biological specimen.

One embodiment of the invention device to collect and immobilize biological specimens is shown in Figure 7. The device 10 is comprised of a matrix 12 for the purpose of immobilizing a biological specimen and a handle 14 to facilitate placement of the matrix 12. The handle 14 is provided to manipulate a small matrix 12 into close proximity of the desired area of the biological specimen and to contact the specimen area and transfer cells 17 from the biological specimen to the matrix 12. The device 10 will further have means to separate the handle 14 from the matrix 12 in a manner that controls placement of the matrix 12 during detachment, allowing the matrix 12 to be placed in a reaction vessel. Quality control manufacturing of the matrix 12 is

expected to maintain consistency in fiber diameter, chemical composition and density. Depending upon the nature of the fibrous matrix, cutting may weaken the ability of the matrix fibers to stay together and require special cutting methods and a backing 16 for physical support. Fabrication of devices 10 comprising very small pieces of matrix material may be facilitated by affixing a length of matrix material to other supporting materials before cutting it into individual pieces for each device.

The embodiment of the device 10 in the drawing of Figure 7 illustrates a design that enables a large assembly of devices 10 to be made at one time. First, a long narrow strip, circa 6 millimeter-width, of matrix material 12 is affixed by means of an adhesive 18 to a plastic backing 16 of the same size to form the matrix sandwich 20.

Handles 14 for a set of devices are fabricated by folding and creasing a sheet of plastic as long as the matrix sandwich 20. A row of openings 22 is made in the sheet so that each handle 14 will have one opening 22. A second adhesive 24 is applied to the first side of the plastic sheet before folding and the matrix sandwich 20 is placed on the second side of the plastic sheet with the backing 16 facing the openings 22. The adhesive 24 fastens the backing 16 of the matrix sandwich 20 to the second side of the plastic sheet through the openings 22. The sheet is then creased at least twice on either side of the row of openings and folded by bringing opposite ends, parallel to the row of openings, and pressing the first side together with the adhesive 24 inside to keep the plastic sheet folded. The matrix sandwich 20 remains affixed to the second side of the plastic sheet by contact with the adhesive 24 through the openings 22. When the assembled sheet is cut perpendicular to the row of openings,

between each pair of openings 20, the matrix strip will also be cut into small pieces in a way that each device has the 12 matrix facing outward as drawn in the single device 10 of Figure 7.

The devices 10 are packaged for individual use and opened at the site of sample collection where a technician will remove the device by its handle 14, guide and contact the matrix 12 to the biological specimen area. To detach the matrix sandwich 20 and place it in a reaction vessel, the handle 14 is unfolded by peeling the first sides apart and away from the adhesive 24, releasing the matrix backing 16 from the adhesive 24 in the area of the opening 22. Adhesives 18 and 24 may be either liquid materials spread over the matrix backing 16 and plastic sheet forming the handles 14, respectively, or a lined, pressure sensitive adhesive transfer tape.

Yet another embodiment of the device comprises a narrow roll of matrix, with or without a backing, contained in a housing having means to advance the roll, cut and release sections of matrix as needed. Housings for applying labels to products and tape to packages are examples of similar mechanisms. Another embodiment of a housing means is similar to a pill dispenser, in which individual matrices are stacked in a cylinder with a first means to expose one matrix at a time for gentle pressure to contact the specimen and a second means to eject the matrix from the container. The invention is a device according to Example 3, further comprising a handle or dispensing apparatus.

It is understood that other embodiments of a sample collecting and processing device are within the scope of this invention, regardless of the mode of manufacturing or the intended type of biological specimen. A preferred embodiment of the device of

the invention are shown as a matrix sandwich comprising nylon and polyester fibers, supported by a plastic backing and a handle from which the matrix may be detached. Different embodiments of a matrix material, upon which the sample is immobilized, include, and are not limited to, fibrous matrices, comprising one or more types of natural or synthetic fibers, that are fibers laid down in a pattern to encourage lateral or vertical flow, absorbent wicking materials, a woven matrix, two or more matrices layered together and the like. The matrix may comprise a sticky surface or a pattern made by manufacturing techniques to increase surface area of a material to which cells and viral particles naturally adhere, or have been coated to increase adhesion. The matrix itself may be an internal part of a reaction vessel. A handle is useful to manipulate and collect a biological specimen, but optional because it is not necessary. The sample collecting matrix, and any other supporting material that becomes part of the reaction, are selected on the basis that they do not inhibit ISA or the analysis of nucleic acids.

A further property of the sample collection device is that it may provide a measure of the amount of biological sample that is collected. For example, a measured amount of a particular fibrous matrix in the device will absorb a predetermined quantity of a liquid biological specimen. Another example of using the device to measure the biological sample is that when the size and adherence of collecting surface area that is provided with the device has been established to collect the same amount of a particular biological specimen within an acceptable range, the results may be expressed in quantifiable terms. Therefore, the invention may be a



method of detecting and determining the relative quantities of nucleic acid targets, such as the prevalence of viral particles, present in a given quantity of specimen.


It is possible to determine the amount of clinical specimen needed for immobilization that will routinely amplify fewer genetic targets than other methods. Immobilizing the sample eliminates extraction procedures and therefore none of the target nucleic acids originally present in the sample are lost due to the fact they are not recovered during the extraction steps. An aqueous solution of biochemically active enzymes and amplification reagents is readily absorbed into a dehydrated matrix and rehydrates the biological specimen. The rehydrated specimen immobilized on the matrix is subjected to appropriate temperature conditions and amplified products in the reaction solution are detected by any number of means known in the art.

In order to detect multiple mutations from the same sample, it is possible to use a high density probe array representing different sequence combinations to which the complementary nucleic acids of the amplified product bind upon diffusing from the immobilized cells. By immobilizing a clinical specimen on a first matrix surface and detection probes on a second surface, both in the same reaction chamber, sample preparation can be integrated with amplification and detection into a single reaction chamber. There are numerous schemes for labeling, transmitting and detecting signals based on the hybridization of complementary nucleic acid sequences that may be applied to detect products of immobilized sample amplification, eliminating the need for gel electrophoresis altogether. The more sensitive the detection, the less amplification product is needed to transmit a signal. The invention is further a method for analyzing the genetic sequence of small amounts of the nucleic acids of a specimen, comprising the

following steps: (a) drying and immobilizing said nucleic acids of said specimen on a first internal surface of a reaction chamber; (b) immobilizing nucleic acids in a known array on a second surface of the reaction chamber; (c) amplifying or labeling said targeted regions of said nucleic acids of said specimen; (d) means for said labeled or amplified nucleic acids of said specimen to bind said nucleic acids immobilized in said known array; (e) and means of detecting the presence, absence or amount of said targeted regions of said nucleic acids of said specimen by interpreting to what genetic sequence of said known array, said labeled or amplified nucleic acids of said specimen are bound by their sequence complementarity.

100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2236  
2237  
2238  
2239  
2240  
2241  
2242  
2243  
2244  
2245  
2246  
2247  
2248  
2249  
2250  
2251  
2252  
2253  
2254  
2255  
2256  
2257  
2258  
2259  
2260  
2261  
2262  
2263  
2264  
2265  
2266  
2267  
2268  
2269  
2270  
2271  
2272  
2273

Applicant (Inventors): Marilyn J. Stapleton Rebecca Sundseth Attorney ocket  
Serial or Patent No.: -- No: STMJ5011  
Filed or Pended: --



VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS  
(37 CFR 1.9(f) AND 1.27(c)) - SMALL BUSINESS CONCERN

I hereby declare that I am

- ☒ the owner of the small business concern identified below:  
☐ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF CONCERN Gene Tec Corporation  
ADDRESS OF CONCERN 205 Winterberry Ridge Drive  
Durham, NC 27713

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.3-18, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled Immobilizing and Processing Specimens on by inventor(s) Marilyn J. Stapleton and  
Rebecca Sundseth described in Matrix Materials for the Identification of  
Nuclear Acid Sequences

- ☒ the specification filed herewith  
☐ application serial no. \_\_\_\_\_, filed \_\_\_\_\_  
☐ patent no. \_\_\_\_\_, issued \_\_\_\_\_

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a non profit organization under 37 CFR 1.9(e). \*Note: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME None  
ADDRESS \_\_\_\_\_  
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date of which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge and after having been warned that willful false statements, and the like, so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF PERSON SIGNING MARILYN J. STAPLETON  
TITLE OF PERSON OTHER THAN OWNER \_\_\_\_\_  
ADDRESS OF PERSON SIGNING \_\_\_\_\_  
205 Winterberry Ridge Drive, Durham

SIGNATURE Marilyn J. Stapleton DATE April 14, 1997

FIG. 1A




FIG. 1B




FIG. 1B

FIG. 2A




FIG. 2B




FIG. 3A





FIG. 3B




FIG. 4




FIG. 5




FIG. 6




FIG. 7

